arXiv:1603.08623v1 [math.NT] 29 Mar 2016

A POSITIVE PROPORTION OF LOCALLY SOLUBLE THUE EQUATIONS ARE GLOBALLY INSOLUBLE

SHABNAM AKHTARI AND MANJUL BHARGAVA

Abstract. We show that a positive proportion of integral binary cubic forms that locally everywhere represent 1 do not globally represent 1. We order all classes of binary cubic forms by their absolute discriminants. We prove the same result for binary forms of any degree $n \geq 3$, provided that these forms are ordered by the maximum of the absolute values of their coefficients.

1. Introduction

The purpose of this article is to prove the existence of many cubic *Thue equations*

$$F(x, y) = 1 \tag{1}$$

that have no solutions (x, y) in the integers; here F is a binary cubic form of nonzero discriminant with coefficients in the integers.

It is easy to construct such forms if we arrange a local obstruction. For example, any binary cubic form congruent to $xy(x+y)$ modulo 2 will never represent an odd number. Also, if $F(x, y) = 1$ has no solution, and G is a *proper subform* of F, i.e., $G(x, y) = F(ax + by, cx + dy)$ for some integer matrix $A = \left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}\right)$ with $|\det A| > 1$, then clearly $G(x, y) = 1$ will also have no integer solutions.

Our goal in this paper is to show for every integer $n \geq 3$ that many (indeed, a positive proportion) of binary forms of degree n are not proper subforms, locally represent 1 at every place, but globally fail to represent 1.

Theorem 1.1. *Let h be any nonzero integer. When $\mathrm{GL}_2(\mathbb{Z})$-classes of integral binary cubic forms $F(x, y) \in \mathbb{Z}[x, y]$ are ordered by absolute discriminant, a positive proportion of these forms F have the following properties:*

(i) *they locally everywhere represent h (i.e., $F(x, y) = h$ has a solution in $\mathbb{R}^2$ and in $\mathbb{Z}_p^2$ for all p);*
(ii) *they globally do not represent h (i.e., $F(x, y) = h$ has no solution in $\mathbb{Z}^2$); and*

Key words and phrases. Binary forms, Thue equations and inequalities, polynomial congruences, Hasse principle.

This work was initiated at the Mathematisches Forschungsinstitut Oberwolfach in summer 2011.

(iii) *they are maximal forms* (*i.e., F is not a proper subform of any other form*).

Our method of proof provides an explicit lower bound for the positive density of classes of binary cubic forms that do not represent a given integer, and indeed an explicit construction of this positive density of forms.

We prove an analogous result for binary forms of general degree, provided that these forms are ordered by the maximum of the absolute values of their coefficients.

Theorem 1.2. *Let h be any nonzero integer. When binary forms $F(x,y) \in \mathbb{Z}[x,y]$ of a given degree $n \geq 3$ are ordered by the maximum of absolute values of their coefficients, a positive proportion of them have the following properties:*

(i) *they locally everywhere represent h (i.e., $F(x,y) = h$ has a solution in $\mathbb{R}^2$ and in $\mathbb{Z}_p^2$ for all p);*
(ii) *they globally do not represent h (i.e., $F(x,y) = h$ has no solution in $\mathbb{Z}^2$); and*
(iii) *they are maximal forms (i.e., F is not a proper subform of any other form).*

As with Theorem 1.1, our method of proof provides an explicit lower bound for the positive density of binary n-ic forms that do not represent a given integer, and moreover, yields an explicit construction of this positive density of forms.

We remark that for every fixed even positive integer n, one can deduce that a positive proportion of binary forms of degree n do not represent a given integer h by using the results in [3], where it is shown that most hyperelliptic curves over $\mathbb{Q}$ have no rational points. However, the methods of [3] do not provide an explicit construction to produce such forms, and also currently apply only in the case that the degree n is even.

2. Preliminaries

Let $F(x,y) \in \mathbb{Z}[x,y]$ be an integral binary form and $A = \left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}\right) \in \mathrm{GL}_2(\mathbb{Q})$. Then define the binary form F_A by

$$F_A(x,y) := F(ax+by,\ cx+dy).$$

If (x_0, y_0) is a solution of $F(x,y) = h$, then

$$A \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \begin{pmatrix} ax_0 + by_0 \\ cx_0 + dy_0 \end{pmatrix}$$

and so $(ax_0 + by_0, cx_0 + dy_0)$ is a solution of $F_{A^{-1}}(x,y) = h$.

If $F(x,y)$ factors in $\mathbb{C}$ as

$$F(x,y) = \prod_{i=1}^{n} (\alpha_i x - \beta_i y),$$

then the discriminant $\mathrm{disc}(F)$ of F is given by

$$\mathrm{disc}(F) = \prod_{i<j}(\alpha_i\beta_j - \alpha_j\beta_i)^2.$$

It follows that, for any 2×2 matrix A, we have

$$\mathrm{disc}(F_A) = (\det A)^{n(n-1)}\mathrm{disc}(F). \tag{2}$$

If $A \in \mathrm{GL}_2(\mathbb{Z})$, then we say that F_A is *equivalent* to F.

A pair $(x_0, y_0) \in \mathbb{Z}^2$ is called a *primitive solution* to the equation $F(x, y) = h$ if $F(x_0, y_0) = h$ and $\gcd(x_0, y_0) = 1$. We denote by $N_{F,h}$ the number of primitive solutions in integers x and y of the equation $F(x, y) = h$. If F_1 and F_2 are equivalent, then evidently $\mathrm{disc}(F_1) = \mathrm{disc}(F_2)$ and $N_{F_1,h} = N_{F_2,h}$.

The following is an immediate consequence of the main results in [1].

Theorem 2.1. *Let $F(x, y) \in \mathbb{Z}[x, y]$ be an irreducible binary form of degree $n \geq 3$ and discriminant D. Let m be an integer with*

$$0 < m \leq \frac{|D|^{\frac{1}{2(n-1)}-\epsilon}}{(3.5)^{n/2}n^{\frac{n}{2(n-1)}}},$$

where $0 < \epsilon < \frac{1}{2(n-1)}$. Then the equation $F(x, y) = m$ has at most

$$\begin{cases} 7n + \frac{n}{(n-1)\epsilon} & \text{if } n \geq 5 \\ 9n + \frac{n}{(n-1)\epsilon} & \text{if } n = 3, 4 \end{cases}$$

primitive solutions. (If n is even, then two solutions (x_0, y_0) and $(-x_0, -y_0)$ are considered here to be one solution).

We will use Theorem 2.1, together with some results on polynomial congruences recalled in the next section, in our construction of binary forms that do not represent 1.

3. Polynomial congruences

In this section, we describe two lemmas on the zeroes of polynomials modulo prime powers, due to Bombieri and Schmidt [6, Lemma 7] and Stewart [14, Theorem 2], which will also form key ingredients in our construction of forms that do not represent 1.

To state these results, let Ω_p be an algebraic closure of the p-adic field $\mathbb{Q}_p$. We denote by $|\ |_p$ the usual absolute value in $\mathbb{Q}_p$, together with an extension to Ω_p. Here $\mathbb{Z}_p$ is the ring of integers of $\mathbb{Q}_p$ and R_p denotes the ring of elements ζ in Ω_p for which $|\zeta|_p \leq 1$. Then we have the following lemma due to Stewart in [14, Theorem 2].

Lemma 3.1. *Let $f(X)$ be a polynomial of degree n with integer coefficients, content 1 and nonzero discriminant D. Let k be a positive integer, p be a prime number, and s be the number of zeros of f in R_p. Then the complete solution of the congruence*

$$f(X) \equiv 0 \ (mod\ p^k)$$

is given by at most s congruences

$$X \equiv c_i \ (mod\ p^{k-u_i}),$$

where c_i are non-negative integers and $0 \leq u_i \leq k$. In addition, the above congruences can be found so that when $u_i \neq 0$, for each root α of $f(X)$, we have

$$|\alpha - c_i|_p \leq p^{u_i - k}. \tag{3}$$

Now suppose $f(X)$ is a polynomial of degree n and content 1. Let p be a prime and k a non-integer. By Lemma 3.1, we can assume that the complete solution of the congruence

$$f(X) \equiv 0 \ (\mathrm{mod}\, p^k)$$

is given by t congruences

$$X \equiv c_i \ (\mathrm{mod}\, p^{k-u_i}),$$

where c_i are non-negative integers and $0 \leq u_i \leq k$. Here t does not exceed the number of zeros of f in R_p and we can order the above congruences so that $u_i > 0$ for $0 < i \leq t_1$ and $u_i = 0$ for $t_1 < i \leq t$.

Set

$$f_i(X) := f(p^{k-u_i}X + b_i).$$

Then we have the following lemma, which is a part of Theorem 2 of [14].

Lemma 3.2. *p^k divides the content of $f_i(X)$, for $i = 1, \ldots t$.*

Thus we can define t (with $t \leq s$) polynomials

$$\tilde{f}_i(X) = p^{-k} f(p^{k-u_i}X + b_i)$$

with integer coefficients.

4. Proof of Theorem 1.1

Our strategy to prove Theorem 1.1 is as follows. We first restrict to the case of $h = 1$, and construct in Subsection 4.1 a set of maximal binary cubic forms $F(x, y)$ of positive density defined by congruence conditions. For any such irreducible binary cubic form $F(x, y)$ of sufficiently large discriminant, we show in Subsection 4.2 that each solution of $F(x, y) = 1$ corresponds uniquely to a solution of $G_j(x, y) = 1$, where G_j lies in a certain set of 81 maximal binary cubic forms $G_j(x, y) = 1$ associated to F with $\mathrm{Disc}(G_j) \ll \mathrm{Disc}(F)$. These G_j's are constructed using the polynomial congruence results of Section 3. Using the fact that F can represent 1 only an absolutely bounded number of times by Theorem 2.1, we conclude in Subsection 4.3 that most of the G_j's cannot represent 1.

In Subsection 4.4, we argue that these forms G_j's that globally do not represent 1 do represent 1 everywhere locally. In Subsection 4.5, we show that the G_j's yield a positive density of all binary cubic forms when ordered by absolute discriminant. This proves Theorem 1.1 in the case $h = 1$. We explain how the proof extends naturally to general h in Subsection 4.6.

We start by assuming $h = 1$.

4.1. **Construction of binary cubic forms $F(x, y)$ satisfying certain congruence conditions and having large discriminant.** Let p_i be the i-th prime greater than 3 and set

$$m = \prod_{i=1}^{k} p_i,$$

where $k \geq 4$ is any integer. Let $F(x, y)$ be a maximal primitive irreducible integral binary cubic form such that

$$|\mathrm{disc}(F)| > m^{20}(3.5)^{30}3^{15}.$$

Assume further that $F(x, y)$ does not split completely modulo 2, does split completely modulo the primes $p_1, \ldots, p_k$, and is irreducible modulo 3. Finally, assume, for each prime $p > p_k$ with $p \equiv 1 \pmod 3$, that $F(x, y)$ does not factor as $cL(x, y)^3$ modulo p for any linear form L and constant c over $\mathbb{F}_p$. We will show in Subsection 4.5 that such forms F form a positive proportion of all classes of binary cubic forms when ordered by absolute discriminant.

4.2. **Construction of forms G_j's associated to each F.** Given any form F as constructed in Subsection 4.1, we will show how to obtain many irreducible maximal binary cubic forms G that locally everywhere represent 1 but do not globally represent 1.

First, because of the condition on the discriminant of F, we have by Theorem 2.1 that the equation

$$F(x, y) = m$$

has at most 34 primitive solutions (take $\epsilon = \frac{1}{5}$).

Assume that

$$F(x, y) \equiv m_0(x - a_1y)(x - b_1y)(x - c_1y) \pmod{p_1}, \tag{4}$$

where $m_0 \not\equiv 0 \pmod{p_1}$, and a_1, b_1 and c_1 are three distinct integers modulo p_1. Following the method of Bombieri and Schmidt in [6], define

$$\begin{aligned} F_{a_1}(x, y) &:= F(p_1x + a_1y, y), \\ F_{b_1}(x, y) &:= F(p_1x + b_1y, y), \\ F_{c_1}(x, y) &:= F(p_1x + c_1y, y). \end{aligned}$$

We claim that these three forms are pairwise inequivalent. One of the entries in the real matrix that transforms $F_{a_1}(x, y)$ to $F_{b_1}(x, y)$ is $\frac{b_1 - a_1}{p_1}$ and since a_1 is not congruent to b_1 modulo p_1, this change of variable matrix does not belong to $\mathrm{GL}_2(\mathbb{Z})$. By Lemma 3.2, or via direct substitution, we see that the contents of $F_{a_1}(x, y)$, $F_{b_1}(x, y)$ and $F_{c_1}(x, y)$ are exactly divisible by p_1. Thus we may divide all the coefficients of F_{a_1}, F_{b_1} and F_{c_1} by p_1 to get three binary forms $\tilde{F}_{a_1}$, $\tilde{F}_{b_1}$ and $\tilde{F}_{c_1}$, with integral coefficients, content 1, and discriminant equal to $p_1^2 \,\mathrm{disc}(F)$.

Since we assumed $F(x,y)$ splits completely modulo p_1, we either have (4) or

$$F(x,y) \equiv m_0 y(x - b_1 y)(x - c_1 y) \pmod{p_1},$$

for some integers $m_0 \not\equiv 0 \pmod{p_1}$, b_1 and c_1, with $b_1 \not\equiv c_1 \pmod{p_1}$. In the latter case, we define F_{b_1} and F_{c_1} as above and set

$$F_\infty(x,y) := F(p_1 y, x)$$

and

$$\tilde{F}_\infty(x,y) := \frac{1}{p_1} F(p_1 y, x).$$

One of the entries in the real matrix that transforms $F_\infty(x,y)$ to $F_{b_1}(x,y)$ is $\frac{1}{p_1}$ and this change of variable matrix does not belong to $\mathrm{GL}_2(\mathbb{Z})$. Therefore, $F_\infty(x,y)$, $F_{b_1}(x,y)$ and $F_{c_1}(x,y)$ are pairwise inequivalent.

By Lemma 3.1, each solution of the equation $F(x,y) = m$ corresponds uniquely to a solution of one of

$$\begin{aligned} \tilde{F}_{a_1}(x,y) &= m/p_1, \\ \tilde{F}_{b_1}(x,y) &= m/p_1, \\ \tilde{F}_{c_1}(x,y) &= m/p_1. \end{aligned}$$

(Here we take $a_1 = \infty$ if $F(x,y)$ is a multiple of y modulo p_1.)

Applying the above procedure, with the prime p_2 in place of p_1, to each of $\tilde{F}_{a_1}(x,y)$, $\tilde{F}_{b_1}(x,y)$ and $\tilde{F}_{c_1}(x,y)$ (note that these forms still split completely modulo $p_2, \dots, p_k$), we then obtain 9 different cubic Thue equations of the shape

$$K(x,y) = \frac{m}{p_1 p_2},$$

where $K(x,y) \in \mathbb{Z}[x,y]$ has discriminant $p_1^2 p_2^2 \mathrm{disc}(F)$.

Applying the above procedure for each prime p_i, $i = 1, \dots, k$, we then obtain 3^k binary cubic forms with integral coefficients, content 1, and discriminant

$$\left(\prod_{i=1}^{k} p_i^2 \right) \mathrm{disc}(F).$$

We denote these binary forms G_j, for $j = 1, \dots, 3^k$.

Note that if $F(x,y)$ is irreducible over $\mathbb{Q}$, then its associated forms $G_j(x,y)$ will also be irreducible over $\mathbb{Q}$. Furthermore, the G_j's are not constructed as proper subforms of the binary form cubic $F(x,y)$, and in fact the G_j's must be maximal forms as well. Indeed, they are maximal over $\mathbb{Z}_p$ for all $p \notin \{p_1, \dots, p_k\}$ (being equivalent, up to a unit constant, to $F(x,y)$ over $\mathbb{Z}_p$ in that case), while for $p \in \{p_1, \dots, p_k\}$, we have $p \nmid \mathrm{disc}(F)$, implying $p^2 || \mathrm{disc}(G_j)$, and so G_j cannot be a subform over $\mathbb{Z}_p$ of any form by equation (2). Hence the G_j's are all maximal forms.

4.3. **Many of the G_j's do not represent 1.** Let F be as in Subsection 4.1, and let the forms G_j, $j = 1, \ldots, 3^k$, be constructed as in Subsection 4.2. Each primitive solution (x_0, y_0) to the cubic equation

$$F(x, y) = m = \prod_{i=1}^{k} p_i$$

corresponds to a unique triple (j, x_j, y_j) so that (x_j, y_j) is a solution to the equation

$$G_j(x, y) = 1.$$

Since the equation $F(x, y) = m$ has at most 34 primitive solutions, at least $3^k - 34$ binary forms among the G_j's cannot represent 1.

Thus starting from any maximal irreducible binary cubic form $F(x, y)$ that splits completely modulo $p_1, p_2, \ldots, p_k$, we have produced at least $3^k - 34$ maximal irreducible binary cubic forms $G(x, y)$ that do not represent 1.

Note that $k = 4$ suffices to produce a positive number of such forms $G(x, y)$: if $k = 4$, then from such an $F(x, y)$ above we obtain $3^4 - 34 = 47$ binary cubic forms $G(x, y)$ such that $G(x, y) = 1$ does not have any solution in integers x and y.

4.4. **The cubic forms F and also the forms G_j's represent 1 everywhere locally.** Next we wish to show that these binary forms G_j that do not globally represent 1 do represent 1 everywhere locally.

Since any binary cubic form takes positive values, we conclude that each G_j certainly takes the value 1 over $\mathbb{R}$. To see that the G_j's represent 1 over $\mathbb{Z}_p$ for all primes p, we use the following lemma.

Lemma 4.1. *A primitive binary cubic form f over $\mathbb{Z}_2$ represents 1 if and only if $f(x, y) \not\equiv xy(x + y) \pmod 2$. If $p \equiv 1 \pmod 6$, then a primitive binary cubic form f over $\mathbb{Z}_p$ represents 1 if and only if $f(x, y) \not\equiv cL(x, y)^3 \pmod p$, where L is a linear form and c is a non-cube modulo p. Finally, if $p \equiv 5 \pmod 6$, then every primitive binary cubic form f over $\mathbb{Z}_p$ represents 1.*

Proof. If $p \equiv 2 \pmod 3$, then every unit in $\mathbb{Z}_p$ is a cube. Thus, in this case, every binary form f over $\mathbb{Z}_p$ that represents a unit represents 1: indeed, if $f(x, y) = u$ for $(x, y) \in \mathbb{Z}_p^2$ and $u \in \mathbb{Z}_p^\times$, then $f(x/a, y/a) = 1$, where $a^3 = u$. Now every primitive binary cubic form over $\mathbb{Z}_p$ represents a unit in $\mathbb{Z}_p$, except when $p = 2$ and $f(x, y) \equiv xy(x + y) \pmod 2$.

It remains to treat the case $p \equiv 1 \pmod 3$. Let c be a cubic non-residue mod p. Then if $f(x, y) \equiv c(x - by)^3$ modulo p, then clearly f does not represent 1 modulo p (for any $b \in \mathbb{F}_p$). We show this is the only case where a local obstruction over $\mathbb{Z}_p$ will arise.

Indeed, if c is a cubic residue modulo p, and $f(x, y) \equiv c(x - by)^3$ modulo p for some b, then f evidently represents unit cubes, and therefore 1, over $\mathbb{Z}_p$.

Next let d, a, and b be three integers, with $d \not\equiv 0$ and $a \not\equiv b \pmod{p}$. Then the binary form

$$d(x-ay)(x-by)^2$$

represents 1 modulo p. This is easy to verify by solving the system of linear equations

$$\begin{aligned} x - ay &= 1 \\ x - by &= d. \end{aligned}$$

Thus if a binary form $f(x,y)$ over $\mathbb{Z}_p$ has a double root but not a triple root modulo p, then it represents a unit cube and therefore 1.

Finally, if a binary form $f(x,y)$ over $\mathbb{Z}_p$ does not have any multiple root modulo p, then the curve

$$z^3 = f(x,y) \pmod{p}$$

over $\mathbb{F}_p$ is smooth of genus 1; therefore by the Hasse bound, the above equation has solutions with $z \not\equiv 0 \pmod{p}$. Thus $f(x,y)$ represents 1 modulo p, so represents unit cubes and thus 1 over $\mathbb{Z}_p$. □

By the assumptions on how F factors modulo each prime, and using the fact that each G_j factors in a predictable way in relation to F modulo every prime, we can then see that each G_j locally represents 1. Indeed, F and thus each G_j is not congruent to $xy(x+y) \pmod{2}$; each $G_j(x,y)$ factors as $c\,L(x,y)y^2$ for some constant c modulo each prime $p \in \{p_1, \ldots, p_k\}$; and each G_j does not factor modulo p as a constant times the cube of a linear form for each prime $p > p_k$ for which $p \equiv 1 \pmod{3}$. By Lemma 4.1, we conclude that F and the G_j's all represent 1 over $\mathbb{Z}_p$ for all $p \neq 3$.

For $p = 3$, we note that the curve $z^3 = G_j(x,y)$ is smooth over $\mathbb{F}_3$, and thus possesses a smooth point over $\mathbb{F}_3$ by the Hasse bound, which then must have nonzero z-coordinate by the irreducibility of G_j. Therefore, $z^3 = G_j(x,y)$ has a solution over $\mathbb{Z}_3$ where z is a unit, and thus $G_j(x,y)$ represents 1 over $\mathbb{Z}_3$ as well. Finally, any nonzero binary cubic form over $\mathbb{R}$ represents positive values and thus represents 1 over $\mathbb{R}$; thus the G_j also all locally represent 1 over $\mathbb{R}$.

In summary, we have the following theorem.

Theorem 4.2. *Let p_i be the i-th odd prime greater than 3 and set $m = \prod_{i=1}^{k} p_i$, where $k \geq 4$ is any integer. Suppose $F(x,y)$ is a maximal primitive irreducible integral binary cubic form, with discriminant greater than $m^{20}(3.5)^{30}3^{15}$, such that $F(x,y)$ is irreducible modulo 3, does not split completely modulo 2, but does split completely modulo the primes $p_1, \ldots, p_k$. Further assume for each prime $p > p_k$ with $p \equiv 1 \pmod{3}$, that $F(x,y)$ does not factor as $cL(x,y)^3$ modulo p for any linear form L and constant c over $\mathbb{F}_p$. Given any such form F, there exist 3^k mutually inequivalent irreducible maximal binary cubic forms G_j, for $j = 1, \ldots, 3^k$, having discriminant $\operatorname{disc}(F)m^2$ such that at least $3^k - 34$ of the G_j satisfy the following:*

(i) *they locally everywhere represent* 1;

(ii) *they globally do not represent* 1; *and*
(iii) *they are maximal forms.*

4.5. **The forms F and also the forms G_j's yield a positive proportion of all classes of forms.** We now verify that the classes of the forms F satisfying the conditions of Theorem 4.2 give a positive proportion of all classes of integral binary cubic forms when ordered by absolute discriminant. For this, we use the following slight refinement of a theorem of Davenport and Heilbronn:

Theorem 4.3 (Davenport–Heilbronn [10])**.** *Let S be a set of integral binary cubic forms defined by congruence conditions modulo bounded powers of primes, where for sufficiently large p: the defining congruence conditions exclude only a set modulo p that is contained in the set of all forms that factor as a constant times the cube of a linear form modulo p. Let $h(S;D)$ denote the number of classes of irreducible binary cubic forms contained in S having discriminant D. Then:*

$$\text{(a)} \quad \sum_{-X<D<0} h(S;D) \sim \frac{\pi^2}{24} X \prod_p \mu_p(S);$$

$$\text{(b)} \quad \sum_{0<D<X} h(S;D) \sim \frac{\pi^2}{72} X \prod_p \mu_p(S),$$

where $\mu_p(S)$ denotes the p-adic density of S in the space of integral binary cubic forms.

Theorem 4.3 states that the number of equivalence classes of irreducible binary cubic forms per discriminant is a positive constant on average, even if we are counting only those forms satisfying a finite (or even a suitable infinite) set of congruence conditions. It is also a theorem of Shintani in [13] that the same is true for reducible forms.

In particular, Theorem 4.3 immediately implies that the set S_0 of all F satisfying the conditions of Theorem 4.2 gives a positive proportion of all classes of integral binary cubic forms, when these classes are ordered by absolute discriminant. Indeed, we may determine the exact density of such F among all irreducible binary cubic forms as follows.

First, we note that if a form is non-maximal, then either it is not primitive, or after an $\mathrm{SL}_2(\mathbb{Z})$-transformation is of the form $a_0x^3+a_1x^2y+a_2xy^2+a_3y^3$, where $p^i \mid a_i$, $i = 0,1,2,3$, for some prime p. Thus for primes $p > p_k$ such that $p \equiv 2 \pmod 3$, the p-adic density of binary cubic forms that are primitive and maximal is given by

$$\mu_p(S_0) = 1 - \frac{1}{p^4} - \frac{p^2-1}{p^6}.$$

The 2-adic density of all binary cubic forms that are primitive, maximal, and not congruent to $xy(x+y) \pmod 2$ is $\mu_2(S_0) = 1 - 1/2^4 - 3/2^6 - 1/2^4 = 53/64$. By [4, Lemma 17], the p_i-adic density (for $i = 1,\dots,k$) of all

primitive forms that split completely modulo p_i is

$$\mu_{p_i} = \frac{1}{6}\frac{(p_i-1)^2 p_i(p_i+1)}{p_i^4};$$

the 3-adic density of all binary cubic forms that are primitive and irreducible is

$$\mu_3 = \frac{1}{3}\frac{(3-1)^2 3(3+1)}{3^4} = 16/81.$$

For primes $p > p_k$ such that $p \equiv 1 \pmod 3$, the density of binary cubic forms that are primitive and not congruent to cL^3 modulo p for some linear form L (and thus automatically maximal) and some constant c over $\mathbb{F}_p$ is given by

$$\mu_p(S_0) = 1 - \frac{(p-1)(p+1)}{p^4}.$$

Therefore, the density of F satisfying the conditions of Theorem 4.2 among all classes of irreducible integral binary cubic forms, when these forms are ordered by absolute discriminant, is given by

$$\frac{53}{64}\cdot\frac{16}{81}\cdot\prod_{i=1}^{k}\frac{1}{6}\frac{p_i^4-p_i^3-p_i^2+p_i}{p_i^4}\prod_{\substack{p>p_k\\ p\equiv 1 \pmod 3}}\left(1-\frac{p^2-1}{p^4}\right)\prod_{\substack{p>p_k\\ p\equiv 2 \pmod 3}}\left(1-\frac{p^2+p+1}{p^6}\right).$$

In order to show that the forms G_j's that we have constructed from these forms F, which are maximal, locally represent 1, but do not globally represent 1, yield a positive proportion of all classes of irreducible integral binary cubic forms, we note that the G_j's are mutually inequivalent, and that each F gives rise to at least 47 G_j's (when $k = 4$), each satisfying

$$\operatorname{disc}(G_j) = \left(\prod_{i=1}^{k} p_i^2\right)\operatorname{disc}(F).$$

We conclude that the density of the classes of our forms G_j among all classes of irreducible integral binary cubic forms is given by

$$47\cdot\frac{53}{64}\cdot\frac{16}{81}\cdot\prod_{i=1}^{k}\frac{1}{6}\frac{p_i^4-p_i^3-p_i^2+p_i}{p_i^4}\frac{1}{p_i^2}\prod_{\substack{p>p_k\\ p\equiv 1 \pmod 3}}\left(1-\frac{p^2-1}{p^4}\right)\prod_{\substack{p>p_k\\ p\equiv 2 \pmod 3}}\left(1-\frac{p^2+p+1}{p^6}\right)$$

when $k = 4$, and we have proven Theorem 1.1 in the case $h = 1$.

4.6. **The case of general h.** We may handle general $h \in \mathbb{Z}$ in a similar manner. Let k again be a positive integer such that $3^k > 34$, let $p_1, \dots, p_k$ be the k smallest primes greater than 3 such that $p_i \nmid h$, and set

$$m = \prod_{i=1}^{k} p_i.$$

Let $F(x, y)$ be a binary cubic form with

$$|\operatorname{disc}(F)| > (hm)^{20}(3.5)^{30}3^{15}.$$

Then by Theorem 2.1, the equation

$$F(x,y) = hm \tag{5}$$

has at most 34 primitive solutions.

In addition to the assumptions on the binary cubic form F that were made in Subsection 4.1, we also assume that for every prime $p \mid h$, we have

$$F(x,y) \equiv L_1(x,y)L_2(x,y)^2 \pmod{p},$$

where $L_1(x,y)$ and $L_2(x,y)$ are linear forms over $\mathbb{F}_p$ that are linearly independent.

Similarly to the case $h = 1$, performing the construction in Lemmas 3.1 and 3.2 as in Subsection 4.2 using the primes $p_1, \dots, p_k$, we obtain 3^k cubic Thue equations of the shape

$$G_j(x,y) = h, \tag{6}$$

so that every solution (x_0, y_0) to the equation (5) corresponds to a solution of one of the equations in (6), for $j \in \{1, \dots, 3^k\}$.

The extra assumption on the factorizationn of F modulo prime factors of h ensures that the equations $F(x,y) = 0$ and $G_j(x,y) = 0$ are solvable over $\mathbb{Z}_p$ for all primes $p \mid h$ as well. Indeed, given $p \mid h$, if $F(x,y) \equiv L_1(x,y)L_2(x,y)^2 \pmod{p}$, then the curve $F(x,y) = h$ over $\mathbb{F}_p$ has a smooth point (namely, on the component $L_1(x,y) = 0$); this smooth point will then lift to a $\mathbb{Z}_p$-point on $F(x,y) = h$. Thus the equations $F(x,y) = h$ and $G_j(x,y) = h$ will again be everywhere locally soluble.

The rest of the proof is identical to the above proof for $h = 1$.

5. Proof of Theorem 1.2

We turn next to Thue equations of general degree. Since asymptotics for the number $\mathrm{GL}_2(\mathbb{Z})$-classes of binary n-ic forms of bounded discriminant are not known for $n > 3$ (see [5]), in this section we instead order all integral binary n-ic forms by the maximum of the absolute values of their coefficients, but otherwise follow a strategy analogous to that of Section 4.

We again begin by restricting first to the case $h = 1$.

5.1. Construction of binary n-ic forms $F(x,y)$ satisfying certain congruence conditions and having large discriminant.

Let p_i be the i-th prime greater than or equal to n and set

$$m = \prod_{i=1}^{k} p_i,$$

where k is a positive integer to be chosen later. Let $F(x,y)$ be a maximal primitive irreducible integral binary form of degree n such that

$$|\mathrm{disc}(F)| > (3.5)^{2n(n-1)}\, n^{2n} \left(\prod_{i=1}^{k} p_i\right)^{4(n-1)}.$$

We assume that the leading coefficient of F is positive, so that F takes positive values. Assume further that $F(x,y)$ splits completely modulo the primes $p_1, \ldots, p_k$. Let $g = (n-1)(n-2)/2$ be the genus of the curve $z^n = F(x,y)$, and assume that for every prime $p \notin \{p_1, \ldots, p_k\}$, with $p < (2g+1)^2$, we have

$$F(x,y) \equiv L_1(x,y)L_2(x,y)^{n-1} \pmod{p^{1+v_p(n)+\delta_2(p,n)}},$$

where $L_1(x,y)$ and $L_2(x,y)$ are linear forms that are linearly independent modulo p, $v_p(n)$ denotes the p-adic valuation of n, and $\delta_2(p,n)$ is equal to 1 if $p = 2$ and $p \mid n$, and is equal to 0 otherwise.

Finally, assume, for each prime $p > (2g+1)^2$, that $F(x,y)$ does not factor as $cL(x,y)^r$ modulo p for any integer $r > 1$ and any linear form $L(x,y)$ and constant c over $\mathbb{F}_p$. We will show in Subsection 5.5 that such forms F form a positive proportion of all classes of binary n-ic forms when ordered by height.

5.2. **Construction of forms G_j's associated to each F.** Assume that

$$F(x,y) \equiv m_0(x - a_1y)(x - a_2y)\ldots(x - a_ny) \pmod{p_1}, \tag{7}$$

where $m_0 \not\equiv 0 \pmod{p_1}$, and $a_1, \ldots, a_n$ are distinct integers modulo p_1. Similarly to the cubic case in Section 4, and following the method of Bombieri and Schmidt [6], for $i = 1, \ldots, n$, define

$$F_{a_i}(x,y) \;:=\; F(p_1x + a_iy, y).$$

We claim that these n forms are pairwise inequivalent. For every $1 \le i < j \le n$, we observe that one of the entries in the real matrix that transforms F_{a_i} to F_{a_j} is $\frac{a_j - a_i}{p}$. Therefore this change of variable matrix does not belong to $\mathrm{GL}_2(\mathbb{Z})$. By Lemma 3.2, the content of $F_{a_i}(x,y)$ is exactly divisible by p_1 for every $i = 1, \ldots, n$. Thus we may divide all the coefficients of F_{a_i}'s by p_1 to get n binary forms $\tilde{F}_{a_i}$, for $i = 1, \ldots, n$, with integral coefficients and content 1. By Lemma 3.1, each solution of the equation $F(x,y) = m$ corresponds uniquely to a solution of one of

$$\tilde{F}_{a_i} = m/p_1, \;\; i = 1, \ldots, n.$$

Since we assumed $F(x,y)$ splits completely modulo p_1, we either have (7) or

$$F(x,y) \equiv m_0y(x - a_2y)\ldots(x - a_ny) \pmod{p_1},$$

for some integers $m_0 \not\equiv 0 \pmod{p_1}$, and distinct $a_2, \ldots, a_n$ modulo p_1. In the latter case, we define $F_{a_2}, \ldots, F_{a_n}$ as above and we take $a_1 = \infty$ and define

$$F_\infty(x,y) := F(p_1y, x)$$

and

$$\tilde{F}_\infty(x,y) := \frac{1}{p_1} F_\infty(x,y).$$

One of the entries of the 2×2 matrix that transforms F_∞ to F_{a_i} is $1/p_1$ and this change of variable matrix does not belong to $\mathrm{GL}_2(\mathbb{Z})$. Therefore, $F_\infty(x, y)$ is not equivalent to any of $F_{a_i}(x, y)$, for $i = 2, \ldots, n$.

In the statement of Theorem 1.2, we order integral binary n-ic forms by their naive heights, where the (*naive*) *height* of an integral binary n-ic form is defined as the maximum of the absolute values of all its coefficients. We now proceed to show that the height of each of the $\tilde{F}_{a_i}$'s can be bounded by a function of the height of F. Let

$$F(x, y) = f_0 x^n + f_1 x^{n-1} y + \ldots + f_n y^n \in \mathbb{Z}[x, y]$$

be a binary form of degree n. Suppose that

$$F(a, 1) \equiv 0 \pmod{p}, \text{ with } a \in \mathbb{Z}. \tag{8}$$

Put

$$F_a(x, y) = F(px + ay, y) = e_0 x^n + e_1 x^{n-1} y + \ldots + e_n y^n. \tag{9}$$

Then

$$e_{n-j} = p^j \sum_{i=0}^{n-j} f_i a^{n-i-j} \binom{n-i}{j}. \tag{10}$$

If $j \geq 1$, clearly e_{n-j} is divisible by p. Since $e_n = F(a, 1)$, by (8), e_n is also divisible by p, as is expected by Lemma 3.2. We also notice that $e_{n-1} = pf'(b)$, where $f'(X)$ denotes the derivative of polynomial $f(X) = F(X, 1)$. Therefore, if a is a simple root modulo p then $f'(b) \not\equiv 0 \pmod{p}$ and

$$\tilde{F}_a(x, y) = \frac{F_a(x, y)}{p}$$

is congruent to

$$y^{n-1} L(x, y) \pmod{p}, \tag{11}$$

where $L(x, y) = l_1 x + l_2 y$ is a linear form modulo p, with $l_1 \not\equiv 0 \pmod{p}$. Lemma 5.1, which will be proven in Subsection 5.4, implies that $\tilde{F}_a(x, y) \equiv 1 \pmod{p}$ is solvable. Further, we may choose the integer a so that

$$-\frac{p-1}{2} \leq a \leq \frac{p-1}{2}. \tag{12}$$

Let $H(F)$ be the naive height of F. Then, by (10), (9) and (12), we have

$$\begin{aligned}
|e_{n-j}| &\leq p^j H(F) \left(\frac{p-1}{2}\right)^{n-j} \sum_{i=0}^{n-j} \left(\frac{p-1}{2}\right)^{-i} \binom{n-i}{j} \\
&\leq p^j H(F) \left(\frac{p-1}{2}\right)^{n-j} \sum_{i=0}^{n-j} \binom{n-i}{j} \\
&= p^j H(F) \left(\frac{p-1}{2}\right)^{n-j} \binom{n+1}{j+1} \\
&\leq p^n H(F) \binom{n+1}{j+1}.
\end{aligned}$$

By Stirling's approximation, for every j, we have

$$\binom{n+1}{j+1} \leq \frac{e\, 2^{n+2}}{2\pi\sqrt{n+1}} < \frac{2^{n+1}}{\sqrt{n+1}}.$$

Therefore, we conclude that

$$H(F_a) < p^n \frac{2^{n+1}}{\sqrt{n+1}}\, H(F),$$

and consequently

$$H(\tilde{F}_a) < p^{n-1} \frac{2^{n+1}}{\sqrt{n+1}}\, H(F). \tag{13}$$

Clearly,

$$H(\tilde{F}_\infty) \leq p^{n-1} H(F),$$

by definition.

Applying the reduction procedure for each prime p_i, $i = 1, \ldots, k$, we obtain n^k binary forms of degree n with integral coefficients, content 1, and naive height bounded above by

$$\left(\prod_{i=1}^{k} p_i\right)^{n-1} \frac{2^{(n+1)k}}{(n+1)^{k/2}}\, H(F).$$

We denote these binary forms G_j for $j = 1, \ldots, n^k$.

If $F(x, y)$ is irreducible over $\mathbb{Q}$ then $G_j(x, y)$ will be irreducible over $\mathbb{Q}$ as well. Finally, the G_j's are not constructed as proper subforms of $F(x, y)$, and so the G_j's must be maximal forms as well. Indeed, they are maximal over $\mathbb{Z}_p$ for all $p \notin \{p_1, \ldots, p_k\}$ (being equivalent, up to a unit constant, to $F(x, y)$ over $\mathbb{Z}_p$ in that case), while for $p \in \{p_1, \ldots, p_k\}$, we have $p \nmid \mathrm{disc}(F)$, implying $p^{(n-1)(n-2)} || \mathrm{disc}(G_j)$, so G_j cannot be a subform over $\mathbb{Z}_p$ of any form by equation (2). Hence the G_j's are all maximal forms.

5.3. **Many of the G_j's do not represent 1.** By Theorem 2.1, because of the condition on the discriminant of F, we conclude that the equation

$$F(x,y) = m$$

has at most $13n$ primitive solutions if $n = 3, 4$, and has at most $11n$ primitive solutions if $n \geq 5$ (take $\epsilon = \frac{1}{4(n-1)}$).

Each primitive solution (x_0, y_0) to the equation

$$F(x,y) = m = \prod_{i=1}^{k} p_i$$

corresponds to a unique triple (j, x_j, y_j) so that (x_j, y_j) is a solution to the equation

$$G_j(x,y) = 1.$$

Since the equation $F(x,y) = m$ has at most $13n$ primitive solutions, at least $n^k - 13n$ binary forms among the G_j's cannot represent 1, if $n = 3, 4$. Similarly, if $n \geq 5$, at least $n^k - 11n$ forms among the G_j's cannot represent 1. Let n be even and assume that a solution (x_0, y_0) to $F(x,y) = m$ corresponds to a solution $(\tilde{x}_0, \tilde{y}_0)$ of $G_I = 1$ for a unique $I \in \{1, \ldots, n^k\}$. Clearly the solution $(-x_0, -y_0)$ to $F(x,y) = m$ will correspond to the solution $(-\tilde{x}_0, -\tilde{y}_0)$ of $G_I = 1$. Hence, as in Theorem 2.1, we may consider (x_0, y_0) and $(-x_0, -y_0)$ as one solution.

Thus starting from any maximal irreducible binary form $F(x,y)$ of degree $n = 3$ or 4 that splits completely modulo $p_1, p_2, \ldots, p_k$, we have produced at least $n^k - 13n$ maximal irreducible binary forms $G(x,y)$ of degree n that do not represent 1. In order to produce a positive number of such forms $G(x,y)$, it suffices to take

$$k > \frac{\log(13n)}{\log n}.$$

Similarly, for $n \geq 5$, we may choose

$$k > \frac{\log(11n)}{\log n}$$

to produce at least $n^k - 11n$ maximal irreducible binary forms $G(x,y)$ of degree n that do not represent 1.

5.4. **The forms G_j's represent 1 everywhere locally.** Next we wish to show that these binary forms $G_j(x,y)$ that do not globally represent 1 do represent 1 everywhere locally.

We first use the following simple lemma to show that the forms G_j's represent 1 modulo every prime $p < (2g+1)^2$.

Lemma 5.1. *Let p be a prime and assume that $F(x,y)$ is a primitive binary n-ic form such that $F(x,y) \equiv L_1(x,y)L_2(x,y)^{n-1} (mod\ p^{1+v_p(n)+\delta_2(p,n)})$, where $L_1(x,y)$ and $L_2(x,y)$ are linearly independent linear forms modulo p. Then F represents 1 over $\mathbb{Z}_p$.*

Proof. We may find (x_0, y_0) such that

$$L_1(x_0, y_0) \equiv L_2(x_0, y_0) \equiv 1 \,(\mathrm{mod}\, p^{1+v_p(n)+\delta_2(p,n)});$$

indeed, since L_1 and L_2 are linearly independent forms modulo p, such $x_0, y_0 \in \mathbb{Z}/p^{1+v_p(n)+\delta_2(p,n)}\mathbb{Z}$ exist. It follows that F represents 1 modulo $p^{1+v_p(n)+\delta_2(p,n)}$. Since a unit in $\mathbb{Z}_p$ is an n-th power if and only if it is an n-th power modulo p^l for sufficiently large l - in fact, $l = 1 + v_p(n) + \delta_2(p, n)$ suffices - we see that F represents unit n-th powers and therefore 1 over $\mathbb{Z}_p$, as desired. □

For any prime number $p < (2g+1)^2$, we see from Lemma 5.1 that each G_j represents 1 in $\mathbb{Z}_p$.

By the Hasse-Weil bound, the number of points N on a curve C of genus g over the finite field $\mathbb{F}_q$ of order q satisfies the inequality

$$|N - (q+1)| \leq 2g\sqrt{q}.$$

For $p > (2g+1)^2$, we assumed that the forms F, and thus the G_j's, do not factor as $c\,L(x,y)^r$ modulo p for any $r > 1$ and any linear form $L(x, y)$ and constant c over $\mathbb{F}_p$; it follows from this assumption that the curves $z^n = F(x, y)$ are irreducible over $\mathbb{F}_p$. The Hasse-Weil bound thus applies to each such curve $z^n = F(x, y)$ to produce a smooth point. Therefore by Hensel's lemma, such a smooth $\mathbb{F}_p$-point will lift to a $\mathbb{Z}_p$-point. We conclude that the forms G_j that we have constructed all represent 1 in $\mathbb{Z}_p$ for primes $p > (2g+1)^2$ as well.

Finally, the G_j's continue to represent positive values and therefore also represent 1 over $\mathbb{R}$. In summary, we have proven the following theorem.

Theorem 5.2. *Let $n \geq 3$ be an integer and let p_i be the i-th prime greater than or equal to n. Set $m = \prod_{i=1}^{k} p_i$ where k is any integer satisfying*

$$\begin{cases} k > \frac{\log(13n)}{\log n} & \text{if } n = 3, 4 \\ k > \frac{\log(11n)}{\log n} & \text{if } n \geq 5. \end{cases}$$

Let $F(x, y)$ be a primitive maximal irreducible integral binary form of degree n with discriminant greater than $(3.5)^{2n(n-1)} n^{2n} m^{4(n-1)}$. Assume further that $F(x, y)$ has positive first coefficient and splits completely modulo the primes $p_1, \ldots, p_k$. Let $g = (n-1)(n-2)/2$ be the genus of the curve $z^n = F(x, y)$ and assume that for every prime p, with $p < p_1$ or $p_k < p < (2g+1)^2$, we have

$$F(x, y) \equiv L_1(x, y) L_2(x, y)^{n-1} \ (\mathit{mod}\ p^{1+v_p(n)+\delta_2(p,n)}),$$

where $L_1(x, y)$ and $L_2(x, y)$ are linearly independent linear forms modulo p. Moreover, assume that for any prime $p > (2g+1)^2$, the form $F(x, y)$ does not factor as $c\,L(x, y)^r$ modulo p for any $r > 1$ and any linear form $L(x, y)$ and constant c over $\mathbb{F}_p$.

Given any such form F, there exist n^k mutually inequivalent irreducible maximal binary cubic forms G_j, for $j = 1, \dots, n^k$, such that at least

$$\begin{cases} n^k - 13n & \text{if } n = 3, 4 \\ n^k - 11n & \text{if } n \geq 5 \end{cases}$$

of the G_j's satisfy the following:

(i) *they locally everywhere represent* 1;
(ii) *they globally do not represent* 1; *and*
(iii) *they are maximal forms.*

5.5. **The forms F and also the forms G_j's yield a positive proportion of all integral binary n-ic forms.** We now verify that the forms F satisfying the conditions of Theorem 5.2 give a positive proportion of all integral binary forms of degree n when ordered by naive height.

Theorem 5.3. *Let S be a set of integral binary n-ic forms defined by congruence conditions modulo bounded powers of primes, where for sufficiently large p: the defining congruence conditions exclude only a set modulo p that is contained in the set of $\mathbb{F}_p$-points of a fixed codimension ≥ 2 subscheme of $\mathbb{A}^{n+1}_{\mathbb{Z}}$. Let $N(S; X)$ denote the number of irreducible binary n-ic forms contained in S having positive leading coefficient and naive height less than X. Then:*

(a) $N(S; X) \sim 2^n X^{n+1} \prod_p \mu_p(S)$;
(b) $N(S; X) \sim 2^n X^{n+1} \prod_p \mu_p(S)$,

where $\mu_p(S)$ denotes the p-adic density of S in the space of integral binary n-ic forms.

Proof. Note that 100% of integral binary n-ic forms, when ordered by naive height, are irreducible by Hilbert's irreducibility theorem. The theorem is then obtained as an immediate consequence of Ekedahl's sieve (see, e.g., [11], [12], or [2] for expositions). □

Theorem 5.3 states that the number of irreducible binary n-ic forms of naive height less than X grows as a constant times X^{n+1}, even if we are counting only those forms satisfying a finite, or even a suitable infinite, set of congruence conditions. In particular, Theorem 5.3 immediately implies that the set S_0 of all F satisfying the conditions of Theorem 5.2 gives a positive proportion of all integral binary n-ic forms, when these forms are ordered by naive height.

Indeed, we may determine the exact density of such F among all irreducible binary n-ic forms as follows. Again, if a form is non-maximal, then either it is not primitive, or after an $\mathrm{SL}_2(\mathbb{Z})$-transformation it is of the form $a_0x^n + a_1x^{n-1}y + \cdots + a_ny^n$, where $p^i \mid a_i$, $i = 0, 1, \dots, n$, for some prime p. In particular, integral binary n-ic forms that are non-maximal must factor modulo some prime p as a constant times the n-th power of a linear form. Thus for primes $p > (2g+1)^2$, the p-adic density of binary n-ic forms that

are primitive and not constant multiples of r-th powers modulo p for any $r > 1$ (and thus are automatically maximal over $\mathbb{Z}_p$) is

$$\Upsilon_p(S_0) = \sum_{r|n} \mu(r) \frac{p^{n/r+1} - 1}{p^{n+1}}, \tag{14}$$

by inclusion–exclusion; here the r-th term corresponds to the density of binary n-ic forms that are constant multiples modulo p of exact r-th powers.

The p_i-adic density (for $i = 1, \dots, k$) of all primitive forms that split completely modulo p_i (and thus are automatically maximal over $\mathbb{Z}_{p_i}$) is

$$\frac{\binom{p_i+1}{n}(p_i - 1)}{p_i^{n+1}}.$$

The p-adic density of primitive binary n-ic forms of the shape

$$L_1(x,y)L_2(x,y)^{n-1} \pmod{p^l} \tag{15}$$

where $L_1(x,y)$ and $L_2(x,y)$ are linearly independent linear forms modulo p and $l = 1 + v_p(n) + \delta_2(p,n)$ (such forms are again automatically maximal over $\mathbb{Z}_p$) is

$$\frac{(p+1)p(p-1)p^{3(l-1)}}{p^{l(n+1)}}.$$

Therefore, the density of F satisfying the conditions of Theorem 5.2 among all classes of irreducible integral binary forms of degree n, when these forms are ordered by height, is at least

$$\frac{1}{2} \cdot \prod_{i=1}^{k} \frac{\binom{p_i+1}{n}(p_i - 1)}{p_i^{n+1}} \prod_{\substack{p \notin \{p_1,\dots,p_k\} \\ p < (2g+1)^2}} \frac{(p^3 - p)p^{3(l-1)}}{p^{l(n+1)}} \prod_{p > (2g+1)^2} \Upsilon_p,$$

where $l = 1 + v_p(n) + \delta_2(p,n)$ depends on p, Υ_p is given as in (14), and $g = (n-1)(n-2)/2$ is the genus of the curve $z^n = F(x,y)$.

In order to show that the forms G_j that we constructed from these forms F, which are maximal, locally represent 1, but do not globally represent 1, yield a positive proportion of all classes of irreducible integral binary n-ic forms, we note that the G_j are mutually inequivalent, and that each F gives rise to $n^k - 11n$ forms G_j (if $n > 4$), each satisfying

$$H(G_j) < \left(\prod_{i=1}^{k} p_i \right)^{n-1} \frac{2^{(n+1)k}}{(n+1)^{k/2}} H(F),$$

by (13). We conclude that the density of our forms G_j, when ordered by height, among all irreducible integral binary n-ic forms is at least

$$(n^k - 13n) \frac{(n+1)^{k/2}}{2^{(n+1)k+1}} \prod_{i=1}^{k} \frac{\binom{p_i+1}{n}(p_i - 1)}{p_i^{2n}} \prod_{\substack{p \notin \{p_1,\dots,p_k\} \\ p < (2g+1)^2}} \frac{(p^3 - p)p^{3(l-1)}}{p^{l(n+1)}} \prod_{p > (2g+1)^2} \Upsilon_p,$$

where again $l = 1 + v_p(n) + \delta_2(p,n)$ depends on p, Υ_p is given in (14) and $g = (n-1)(n-2)/2$ is the genus of the curve $z^n = F(x,y)$, and $n = 3, 4$.

Similarly, if $n \geq 5$, we conclude that the density of our forms G_j, when ordered by height, among all irreducible integral binary n-ic forms is at least

$$(n^k - 11n)\frac{(n+1)^{k/2}}{2^{(n+1)k+1}} \prod_{i=1}^{k} \frac{\binom{p_i+1}{n}(p_i-1)}{p_i^{2n}} \prod_{\substack{p \notin \{p_1,\ldots,p_k\} \\ p < (2g+1)^2}} \frac{(p^3-p)p^{3(l-1)}}{p^{l(n+1)}} \prod_{p > (2g+1)^2} \Upsilon_p.$$

5.6. **The case of general h.** We may handle general $h \in \mathbb{Z}$ in a similar manner. Let k be a positive integer satisfying

$$\begin{cases} k > \frac{\log(13n)}{\log n} & \text{if } n = 3, 4 \\ k > \frac{\log(11n)}{\log n} & \text{if } n \geq 5. \end{cases}$$

Let $p_1, \ldots, p_k$ be the k smallest primes that are greater than or equal to n such that $p_i \nmid h$, and set $m = \prod_{i=1}^{k} p_i$.

Let $F(x,y)$ be an irreducible n-ic binary form, with

$$|\mathrm{disc}(F)| > (3.5)^{2n(n-1)} n^{2n} (hm)^{4(n-1)}.$$

Then by Theorem 2.1, the equation

$$F(x,y) = hm \tag{16}$$

has at most $13n$ primitive solutions if $n = 3, 4$, and has at most $11n$ primitive solutions if $n \geq 5$.

In addition to the assumptions on the form F that were made in Subsection 5.1, we also assume that for every prime $p \mid h$, we have

$$F(x,y) \equiv L_1(x,y) L_2(x,y)^{n-1} \pmod{p},$$

where $L_1(x,y)$ and $L_2(x,y)$ are linear forms over $\mathbb{F}_p$ that are linearly independent.

Similarly to the case $h = 1$, performing the construction in Lemmas 3.1 and 3.2 as in Subsection 5.2 using the primes $p_1, \ldots, p_k$, we obtain n^k Thue equations of the shape

$$G_j(x,y) = h, \tag{17}$$

so that every solution (x_0, y_0) to the equation (16) corresponds to a solution of one of the equations in (17), for $j \in \{1, \ldots, n^k\}$.

The extra assumption on the factorization of F modulo prime factors of h again ensures, as in Subsection 4.6 in the case $n = 3$, that the equations $F(x,y) = 0$ and $G_j(x,y) = 0$ are solvable over $\mathbb{Z}_p$ for all primes $p \mid h$ as well.

The rest of the proof is identical to the above proof for $h = 1$.

References

[1] S. Akhtari, *Representation of small integers by binary forms*, Q. J. Math 66 (4) (2015), 1009-1054.

[2] M. Bhargava, *The geometric sieve and the density of squarefree values of invariant polynomials*, arXiv:1402.0031.

[3] M. Bhargava, *Most hyperelliptic curves over $\mathbb{Q}$ have no rational points*, arXiv:1308.0395v1.

[4] M. Bhargava, A. Shankar and J. Tsimerman, *On the Davenport-Heilbronn theorem and second order terms*, Invent. Math. 193 (2013), no. 2, 439-499.

[5] B.J. Birch and J.R. Merriman, *Finiteness theorems for binary forms with given discriminant*, Proc. London Math. Soc. 25 (1972), 385-394.

[6] E. Bombieri, W. M. Schmidt, *On Thue's equation*, Invent. Math. 88 (1987), 69-81.

[7] B.N. Delone and D. K. Faddeev, *The theory of irrationalities of the third degree*, AMS Translations of Mathematical Monographs 10, 1964.

[8] H. Davenport, *On a principle of Lipschitz*, J. London Math. Soc. 26 (1951), 179-183. Corrigendum: On a principle of Lipschitz, J. London Math. Soc. 39 (1964), 580.

[9] H. Davenport, *On the class-number of binary cubic forms I and II*, J. London Math. Soc. 26 (1951), 183-198.

[10] H. Davenport and H. Heilbronn, *On the density of discriminants of cubic fields II*, Proc. Roy. Soc. London Ser. A 322 (1971), no. 1551, 405-420.

[11] T. Ekedahl , *An infinite version of the Chinese remainder theorem*, Comment. Math. Univ. St. Paul. 40 (1991), no. 1, 53-59.

[12] B. Poonen, M. Stoll, *A local-global principle for densities*, in: Scott D. Ahlgren (ed.) et al.: Topics in number theory. In honor of B. Gordon and S. Chowla. Kluwer Academic Publishers, Dordrecht. Math. Appl., Dordr. 467 (1999), 241-244.

[13] T. Shintani, *On Dirichlet series whose coefficients are class numbers of integral binary cubic forms*, J. Math. Soc. Japan 24, Number 1 (1972), 132-188.

[14] C. L. Stewart, *On the number of solutions of polynomial congruences and Thue equations*, Journal of American Math. Soc. 4 (1991), 793-838.

Fenton Hall, University of Oregon, Eugene, OR 97403-1222 USA
E-mail address: akhtari@uoregon.edu

Fine Hall, Washington Road, Princeton NJ 08544-1000 USA
E-mail address: bhargava@math.princeton.edu